

COMPARISON OF NEUTRINO AND MUON STRUCTURE FUNCTIONS, SHADOWING CORRECTIONS AND CHARGE SYMMETRY VIOLATION

C. BOROS AND A. W. THOMAS

*Department of Physics and Mathematical Physics, and Special Research Center for the Subatomic Structure of Matter,
University of Adelaide, Adelaide 5005, Australia*

J. T. LONDERGAN

Department of Physics and Nuclear Theory Center, Indiana University, Bloomington, IN 47404, USA

Comparison of structure functions measured in neutrino and charged lepton deep inelastic scattering can be used to test basic properties of parton distribution functions such as the validity of charge symmetry. Recent experiments indicate a substantial discrepancy between F_2^ν and F_2^μ in the region of small Bjorken- x . We discuss nuclear corrections and strange and anti strange quark effects and show that none of them can account for the observed discrepancy. These results suggest surprisingly large CSV effects in nucleon sea distributions.

1 Introduction

Comparison of structure functions measured in neutrino and charged lepton deep inelastic scattering can be used to test basic symmetry properties of parton distribution functions such as the validity of charge symmetry¹.

Such comparisons are based on the interpretation of these structure functions in terms of parton distribution functions. In the quark-parton model the structure functions measured in neutrino, anti neutrino and charged lepton DIS on an iso-scalar target, N_0 , are given by²

$$\begin{aligned} F_2^{\nu N_0}(x) &= x[Q(x) + 2s(x) - \delta u(x) - \delta \bar{d}(x)] \\ F_2^{\bar{\nu} N_0}(x) &= x[Q(x) + 2\bar{s}(x) - \delta d(x) - \delta \bar{u}(x)] \\ F_2^{\ell N_0}(x) &= \frac{5}{18}x[Q(x) + \frac{2}{5}(s(x) + \bar{s}(x)) \\ &\quad - \frac{4(\delta d(x) + \delta \bar{d}(x))}{5} - \frac{(\delta u(x) + \delta \bar{u}(x))}{5}] \quad (1) \end{aligned}$$

Here, we introduce the notation $Q(x) = u(x) + \bar{u}(x) + d(x) + \bar{d}(x)$ and express everything in terms of parton distributions functions in the proton and charge symmetry violating distributions which are defined as the differences between the up (down) quark distribution in the proton and the down (up) quark distribution in the neutron

$$\begin{aligned} \delta u(x) &= u^p(x) - d^n(x) \\ \delta d(x) &= d^p(x) - u^n(x). \quad (2) \end{aligned}$$

If charge symmetry is valid these terms are zero.

A very useful quantity in the comparison of the structure functions is the “charge ratio”

$$R_c(x) \equiv \frac{F_2^{\mu N_0}(x)}{\frac{5}{18}F_2^{\nu N_0}(x) - x[s(x) + \bar{s}(x)]/6}. \quad (3)$$

A deviation $R_c(x) \neq 1$, at any value of x , must arise either from CSV effects or from $s(x) \neq \bar{s}(x)$.

Recent experimental measurements allow a precise comparison between $F_2^\nu(x, Q^2)$ and $F_2^\mu(x, Q^2)$. In Fig.1 we show the “charge ratio” calculated by using the CCFR³ and NMC⁴ structure functions for F_2^ν and F_2^μ , respectively. The structure functions are integrated over $Q^2 = 3.2 \text{ GeV}^2$ in the region of overlap of the two experiments. Since the CCFR structure function has been measured on an iron target nuclear corrections (nuclear EMC effect, shadowing and anti shadowing) have to be applied to the data. In Fig. 1, the open triangles represent the result without nuclear corrections. The open circles are calculated by using nuclear corrections obtained from *charged lepton* DIS. While, in the region of intermediate values of Bjorken x ($0.1 \leq x \leq 0.4$), the two structure functions are in very good agreement, in the small x -region ($x < 0.1$), they differ by as much as 10-15% when nuclear corrections are taken into account. This discrepancy could be interpreted as evidence for charge symmetry violation. However, several corrections have to be applied to the data before any conclusions may be drawn. We see especially that the result is very sensitive to nuclear corrections. The CCFR Collaboration made a careful study of overall normalization, charm threshold and iso-scalar correction effects³. Here, we re-examine nuclear corrections for neutrinos and discuss $s(x) \neq \bar{s}(x)$ effects before we turn to possible charge symmetry violation.

2 Nuclear corrections

Nuclear corrections for neutrinos are generally calculated using correction factors from charged lepton reactions at the same kinematic values. *A priori*, there is no reason that neutrino and charged lepton heavy target correc-




Figure 1: The “charge ratio” R_c vs. x calculated using CCFR³ data for neutrino and NMC⁴ data for muon structure functions. Open triangles: no heavy target corrections; open circles: ν data corrected for heavy target effects using corrections from charged lepton scattering; solid circles: ν shadowing corrections calculated in the “two phase” model. Both statistical and systematic errors are shown.

tions should be identical, especially if such corrections depend strongly on the properties of the exchanged object (photon, W) used to probe the structure of the target. Since this is the case for nuclear shadowing corrections in the small x_B region for small to moderately large Q^2 -values we re-examined shadowing corrections to neutrino DIS focusing on the differences between neutrino and charge lepton scattering and on effects due to the Q^2 -dependence of shadowing. We used a two phase model which has been successfully applied to the description of shadowing in charged lepton DIS⁵. In generalizing this approach to weak currents, subtle differences between shadowing in neutrino and charged lepton DIS arise because of the partial conservation of axial currents (PCAC) and the coupling of the weak current to both vector and axial vector mesons. PCAC requires that that the divergence of the axial current does not vanish but is proportional to the pion field for $Q^2 = 0$. This is Adler’s theorem⁶, which relates the neutrino cross section to the pion cross section on the same target for $Q^2 = 0$. Thus, for low $Q^2 \approx m_\pi^2$ shadowing in neutrino scattering is determined by the absorption of pions on the target.

For larger Q^2 -values the contributions of vector and axial vector mesons become important. The coupling of the weak current to the vector and axial vector mesons and that of the electro-magnetic current to vector mesons are related to each other by the “Weinberg sum rule” $f_{\rho^+}^2 = f_{a_1}^2 = 2f_{\rho^0}^2$. Since the coupling of the vector (axial vector) mesons to the weak current is twice as large as the coupling to the electro-magnetic current but the structure function is larger by a factor of $\sim 18/5$ in the

neutrino case, we expect that shadowing due to VMD in neutrino reactions is roughly half of that in charged lepton scattering. For large Q^2 -values, shadowing due to Pomeron exchange (which is of leading twist) becomes dominant, leading to identical (relative) shadowing in neutrino and charged lepton DIS.

We calculated shadowing corrections⁷ to the CCFR ν structure function in the framework of this “two phase” model and used these corrections in calculating the charge ratio R_c of Eq. 3. There are also nuclear effects in the Deuteron. However, because of the low density of the Deuteron, these are (relatively speaking) very small and have a negligible effect on the charge ratio. We integrated the structure functions above $Q^2 = 3.2 \text{ GeV}^2$ in the overlapping kinematic region of the two experiments and used a parametrization of the nuclear corrections in charged lepton DIS to correct the data in the non-shadowing region. The result is shown as solid circles in Fig.1. At small x , careful consideration of neutrino shadowing corrections decreases, but does not resolve, the low- x discrepancy between the CCFR and NMC data.

3 Strange and anti strange quark effects

Since the CCFR-Collaboration uses both neutrino and anti neutrino events in the structure function analysis the extracted structure function F_2^{CCFR} is a flux weighted average between ν and $\bar{\nu}$ structure functions³

$$F_2^{CCFR}(x, Q^2) = \alpha F_2^\nu(x, Q^2) + (1 - \alpha) F_2^{\bar{\nu}}(x, Q^2). \quad (4)$$

Here, we define the relative neutrino flux as $\alpha = \Phi_\nu / (\Phi_\nu + \Phi_{\bar{\nu}})$, where Φ_ν and $\Phi_{\bar{\nu}}$ are the ν and $\bar{\nu}$ fluxes, respectively. F_2^{CCFR} is equal to $\frac{1}{2}[F_2^\nu(x, Q^2) + F_2^{\bar{\nu}}(x, Q^2)]$ if $\alpha = \frac{1}{2}$ or if the two structure functions are equal, i.e. charge symmetry is valid and $s(x) = \bar{s}(x)$. The value of α depends on the energy of the incident neutrinos and anti neutrinos in the CCFR-experiment. In the relevant kinematic region, at small x which corresponds to large incident energies, it is ≈ 0.83 so to a good approximation $F_2^{CCFR}(x, Q^2)$ can be regarded as a neutrino structure function.

The remaining small- x discrepancy in the charge ratio is either from different strange quark distributions $s(x) \neq \bar{s}(x)$ or from charge symmetry violation. First, we examine the role played by the strange quark distributions. Assuming charge symmetry, $s(x)$ and $\bar{s}(x)$ are given by a linear combination of neutrino and muon structure functions,

$$\begin{aligned} \frac{5}{6} F_2^{CCFR}(x, Q^2) - 3 F_2^{NMC}(x, Q^2) &= \frac{1}{2} x [s(x) + \bar{s}(x)] \\ &+ \frac{5}{6} (2\alpha - 1) x [s(x) - \bar{s}(x)]. \end{aligned} \quad (5)$$




Figure 2: $xs(x)$ (open circles) and $x\bar{s}(x)$ (solid circles) extracted by combining CCFR and NMC structure functions with $s(x)^{\mu\mu}$ extracted from dimuon experiments, as given in Eqs. 5 and 6. Solid triangles: $\frac{5}{6}F_2^{CCFR} - 3F_2^{NMC}$. Stars: $xs(x)$ from a LO-analysis⁸. Solid line: $xs(x)$ from a NLO-analysis⁹; dashed band indicates $\pm 1\sigma$ uncertainty.

Under the assumption $s(x) = \bar{s}(x)$, this relation could be used to extract the strange quark distribution. However, as is well known, $s(x)$ obtained in this way is inconsistent with results extracted from independent experiments.

Opposite sign dimuon production in deep inelastic ν and $\bar{\nu}$ scattering provides a direct determination of both $s(x)$ and $\bar{s}(x)$. The CCFR Collaboration extracted $s(x)$ and $\bar{s}(x)$ from a leading order (LO)⁸ next to leading order (NLO) analysis⁹ of their dimuon data. While the strange and anti strange distributions were different in the LO analysis they were equal within experimental errors in the NLO analysis.

To test the hypothesis that the low- x discrepancy in the charge ratio of Eq. 3 could be accounted for by allowing $s(x) \neq \bar{s}(x)$ we combined the dimuon production data, averaged over ν and $\bar{\nu}$ events, with the structure functions from neutrino and charged lepton scattering (Eq. 5). Defining $\alpha' = N_\nu / (N_\nu + N_{\bar{\nu}})$, where $N_\nu = 5,030$, $N_{\bar{\nu}} = 1,060$ ($\alpha' \approx 0.83$) are respectively the ν and $\bar{\nu}$ events from the dimuon production experiment⁹, the flux-weighted experimental distribution $xs(x)^{\mu\mu}$ from dimuon production is

$$xs^{\mu\mu}(x) = \frac{1}{2}x[s(x) + \bar{s}(x)] + \frac{1}{2}(2\alpha' - 1)x[s(x) - \bar{s}(x)]. \quad (6)$$

Eqs. (5) and (6) form a pair of linear equations which can be solved for $s(x)$ and $\bar{s}(x)$. We can simultaneously test the compatibility of the various experiments.

Compatibility of the two experiments requires that physically acceptable solutions for $\frac{1}{2}x[s(x) + \bar{s}(x)]$ and

$\frac{1}{2}x[s(x) - \bar{s}(x)]$, satisfying both Eq. 5 and Eq. 6, can be found. Clearly, solutions do not exist if $\frac{5}{3}(2\alpha - 1) = (2\alpha' - 1)$. (Note that the left-hand sides of Eq. 5 and Eq. 6 are different.) This gives the critical values $\alpha = \frac{1}{5}(3\alpha' + 1)$ or $\alpha \approx 0.7$ for $\alpha' \approx 0.83$ and $\alpha = 0.8$ for $\alpha' = 1$.

In Fig. 2 we show the results obtained for $xs(x)$ (open circles) and $x\bar{s}(x)$ (solid circles) by solving the resulting linear equations, Eqs. 5 and 6, with $\alpha \approx \alpha' \approx 0.83$. Both the structure functions and dimuon data have been integrated over $Q^2 > 3.2 \text{ GeV}^2$ in the overlapping kinematical regions. In averaging the dimuon data we used the CTEQ4L parametrization for $s^{\mu\mu}(x)$ ¹⁰. We see that the results are completely unphysical, since the equations require $\bar{s}(x) < 0$. Since $\alpha \approx 0.83$ is closer to the critical value for $\alpha' = 1$ than for $\alpha' = 0.83$ the results is even more unphysical if we use only the neutrino dimuon data. Similarly, if we decrease α $\bar{s}(x)$ becomes more negative.

In conclusion, our analysis strongly suggests that requiring charge symmetry, but allowing $s(x) \neq \bar{s}(x)$, cannot resolve the discrepancy between $F_2^{CCFR}(x, Q^2)$ and $F_2^{NMC}(x, Q^2)$. The experimental results are incompatible, even if $\bar{s}(x)$ is completely unconstrained.

4 Charge symmetry violation

We have seen that neither neutrino shadowing corrections nor allowing $s(x) \neq \bar{s}(x)$ removes the low- x discrepancy. There remain the two possible explanations. Either one of the experimental structure functions (or $s(x)$) is incorrect at low x , or parton charge symmetry is violated in this region. Assuming the possibility of parton CSV, we can combine the dimuon data for $s(x)$, (Eq. 6), with Eq. 5 to obtain

$$\begin{aligned} & \frac{5}{6}F_2^{CCFR}(x, Q^2) - 3F_2^{NMC}(x, Q^2) - xs^{\mu\mu}(x) \\ & \approx \frac{1}{2}[\frac{5}{3}(2\alpha - 1) - (2\alpha' - 1)]x[s(x) - \bar{s}(x)] \\ & \quad + \frac{1}{2}x[\delta\bar{q}(x) - \delta\bar{u}(x)]. \end{aligned} \quad (7)$$

This equation is valid at small x , where sea quark distributions are much larger than valence quarks, so we make the simplest assumption, namely that $\delta q_v(x) = \delta q(x) - \delta\bar{q}(x) \approx 0$.

The left hand side of Eq. 7 and the coefficient in front of $s - \bar{s}$ are positive. Consequently, the smallest CSV effects will be obtained when $\bar{s}(x) = 0$. Note that this violates the requirement that the net strangeness of the nucleons be zero. Thus, setting $\bar{s}(x) = 0$ gives the absolute lower limit on charge symmetry violation. In Fig. 3 we show the CSV effects needed to satisfy the experimental values in Eq. 7. The open circles are obtained when we set $\bar{s}(x) = 0$, and the solid circles result from




Figure 3: Charge symmetry violating distributions extracted from the CCFR and NMC structure function data and the CCFR dimuon production data under the assumption that $s(x) = \bar{s}(x)$ (solid circles) and $\bar{s}(x) \approx 0$ (open circles) for $\alpha' = 0.83$ and $s(x) = \bar{s}(x)$ (solid circles) and $\bar{s}(x) \approx 0$ (open triangles) for $\alpha' = 1$. (For the latter only statistical errors are shown.) $s(x)$ at $Q^2 = 4$ GeV² obtained by the CCFR Collaboration in a NLO analysis⁹ is shown for comparison.

setting $\bar{s}(x) = s(x)$. We also show the results we obtain by setting $\alpha' = 1$ which corresponds to using only a subsample of the di-muon data containing only neutrino events. For $\bar{s}(x) = s(x)$, the result is the same. However, setting $\bar{s}(x) = 0$ gives us a higher lower limit on CSV than that for α' . The result is shown as open triangles in Fig. 3.

The magnitude of the extracted CSV is also sensitive to the strange quark distribution used in the analysis. In Fig. 4 we show the CSV obtained by using the LO (open circles) and NLO (solid triangles) CCFR distributions and the CTEQ4L (solid circles) and CTEQ4D (solid rectangles) parametrizations. The uncertainty due to different parametrizations has been partly taken into account since the calculated errors already include the uncertainty of the dimuon measurement and most of the parametrizations lie within the experimental errors of the dimuon data (except of LO-CCFR $s(x)$).

In conclusion, the CSV effect required to account for the low- x NMC-CCFR discrepancy is extraordinarily large. It is roughly the same size as the strange quark distribution at small x . This CSV term is roughly 25% of the light sea quark distributions for $x < 0.1$, and the sign gives $\bar{d}^p(x) > \bar{u}^n(x)$ and $\bar{d}^n(x) > \bar{u}^p(x)$.

5 Influence of CSV on other observables

Clearly, CSV effects of this magnitude need further experimental verification. It is hard to imagine how such large CSV effects are compatible with the high precision




Figure 4: Uncertainty due to the parametrization used for the dimuon data on charge symmetry violation. Open circles: LO CCFR distribution, solid circles: CTEQ4L, solid rectangles: CTEQ4D, solid triangles: NLO CCFR distribution. Here, except for the most “critical” point, only statistical errors are shown.

of charge symmetry measured at low energies. The level of CSV required is surprising, as it is at least two orders of magnitude larger than theoretical CSV estimates^{11,12}. Theoretical considerations suggest that $\delta\bar{d}(x) \approx -\delta\bar{u}(x)$ ¹²; with this sign CSV effects also require large flavor symmetry violation. Since most of the observables are proportional to the sum of $\delta\bar{d}(x)$ and $\delta\bar{u}(x)$ rather than to their difference this large CSV could remain unobserved in many experiments. Here, we briefly discuss the effects of the extracted CSV on FSV.

The Drell-Yan experiment measures the ratio of the dimuon cross sections of proton-Deuteron and proton-proton scattering. Since CSV is significant in the small x region, it is a good approximation to keep only the contributions to the Drell-Yan cross sections which come from the annihilation of quarks of the projectile and anti quarks of the target. In this approximation, the ratio $R \equiv \sigma^{pD}/(2\sigma^{pp})$ is given by

$$\frac{\sigma^{pD}}{2\sigma^{pp}} \approx \frac{1}{2} \frac{[1 + \frac{\bar{d}_2}{\bar{u}_2} - \frac{\delta\bar{d}_2}{\bar{u}_2}] + \frac{R_1}{4}[1 + \frac{\bar{d}_2}{\bar{u}_2} - \frac{\delta\bar{u}_2}{\bar{u}_2}]}{1 + \frac{R_1}{4} \frac{\bar{d}_2}{\bar{u}_2}}. \quad (8)$$

Here, we introduced the notation $R_1 \equiv \frac{q_1}{q_2}$ and $q_{1,2} \equiv q(x_{1,2})$ for the quark distributions. Neglecting R_1 for large x_F , which corresponds to large x_1 , we have

$$R = \frac{\sigma^{pD}}{2\sigma^{pp}} \approx \frac{1}{2} \{1 + \frac{(\bar{d}_2 - \delta\bar{d}_2)}{\bar{u}_2}\}. \quad (9)$$

If charge symmetry is violated, the extracted quantity¹³ is not $\bar{d}_2/\bar{u}_2$ but $r_2 \equiv (\bar{d}_2 - \delta\bar{d}_2)/\bar{u}_2$. Since $\delta\bar{d}$ is positive the FSV ratio, $\bar{d}_2/\bar{u}_2$, should be enhanced for small x relative to the measured value r_2 . The enhancement is




Figure 5: The ratio $\bar{d}/\bar{u}$ extracted from the Drell-Yan data assuming the validity of charge symmetry (solid circles). If CS is violated this ratio corresponds to $(\bar{d} - \delta\bar{d})/\bar{u}$. The result obtained by correcting for CSV is shown as open circles. The same for the ratio extracted from the difference of the proton and Deuteron structure functions is shown as solid and open triangles, with and without CSV, respectively.

in the order of 25% in the small x region where CSV could be important. This is shown in Fig. 5.

The difference, $\bar{d} - \bar{u}$, can also be extracted from the difference between the proton and neutron structure functions measured by the the NMC Collaboration⁴. In this case we have

$$\frac{1}{2}(u_v - d_v) - \frac{3}{2x}(F_2^p - F_2^n) = (\bar{d} - \bar{u}) - \frac{2}{3}(\delta d + \delta\bar{d}) - \frac{1}{6}(\delta u + \delta\bar{u}). \quad (10)$$

We can make the approximations $\delta q \approx \delta\bar{q}$ and $\delta\bar{d} \approx -\delta\bar{u}$, and obtain

$$\frac{1}{2}(u_v - d_v) - \frac{3}{2x}(F_2^p - F_2^n) \approx [(\bar{d} - \delta\bar{d}) - \bar{u}]. \quad (11)$$

We see that, in a first approximation, the quantities extracted from the two experiments are the same even if both CSV and FSV are present. However, if CSV is present, the term $\delta\bar{d}$ has to be subtracted from the measured quantity to obtain the difference $\bar{d} - \bar{u}$.

We inverted Eq. 11 by dividing both sides by $\bar{d} - \delta\bar{d} + \bar{u} \equiv \bar{u}(r_2 + 1)$, approximating $\bar{d} - \delta\bar{d} + \bar{u}$ on the left hand side of Eq. 11 by a parametrization of $\bar{d} + \bar{u}$ and solving for r_2 . The structure functions and the parton distribution are integrated for each data point over the same Q^2 regions as in the analysis of the charge ratio. The result is shown in Fig. 5 as solid triangles. If we subtract the contribution of CSV from the ratio r_2 we obtain the result shown as open triangles in Fig. 5.

6 Conclusions

In conclusion, we have examined in detail the discrepancy at small x between the CCFR neutrino and NMC muon structure functions. The only way we can make these data compatible is by assuming charge symmetry violation in the sea quark distributions. The CSV amplitudes necessary to obtain agreement with experiment are extremely large – at least two orders of magnitude greater than theoretical predictions of charge symmetry violation. If CSV effects of this magnitude are really present, then one must include charge symmetry violating quark distributions in phenomenological models from the outset, and re-analyze the extraction of parton distributions.

Acknowledgements

This work is supported in part by the Australian Research Council, and by the National Science Foundation under research contract nsf-phy9722706.

References

1. G. A. Miller, B. M. K. Nefkens and I. Slaus, *Phys. Rep.* **194** (1990) 1; E. M. Henley and G. A. Miller in *Mesons in Nuclei*, eds M. Rho and D. H. Wilkinson (North-Holland, Amsterdam 1979).
2. J.T. Londergan and A.W. Thomas, to be published in *Progress in Particle and Nuclear Physics*, 1998.
3. CCFR-Collaboration, W.G.Seligman *et al.*, *Phys. Rev. Lett.* **79**, 1213 (1997).
4. New Muon Collaboration (NMC), M.Arneodo *et al.*, *Nucl. Phys.* **B483**, 3 (1997).
5. J. Kwiecinski and B. Badelek, *Phys. Lett.* **B208**, 508 (1988); W. Melnitchouk and A.W. Thomas, *Phys. Lett.* **B317**, 437 (1993); *Phys. Rev.* **C52**, 3373 (1995).
6. S. L. Adler *Phys. Rev.* **B135**, 963 (1964).
7. C. Boros, J. T. Londergan and A. W. Thomas, (preprint hep-ph/9804411).
8. S.A.Rabinowitz *et al.*, CCFR-Collaboration, *Phys. Rev. Lett.* **70**, 134 (1993).
9. CCFR-Collaboration, A.O. Bazarko *et al.*, *Z. Phys.* **C65**, 189 (1995).
10. H. L. Lai *et al.*, *Phys. Rev.* **D55**, 1280 (1997).
11. E. Sather, *Phys. Lett.* **B 274**, 433 (1992); C.J. Benesh and T. Goldman, *Phys. Rev.* **C55**, 441 (1997).
12. E. Rodionov, A. W. Thomas and J. T. Londergan, *Mod. Phys. Lett.* **A9**, 1799 (1994); C.J. Benesh and J.T. Londergan, preprint *nucl-th/9803017*.
13. E866-Collaboration, E. A. Hawker *et al.*, *Phys. Rev. Lett.* **80**, 3715 (1998)